

SCHUBERT POLYNOMIALS AND QUIVER FORMULAS

ANDERS S. BUCH, ANDREW KRESCH, HARRY TAMVAKIS, AND ALEXANDER YONG

1. INTRODUCTION

The work of Buch and Fulton [BF] established a formula for a general kind of degeneracy locus associated to an oriented quiver of type A . The main ingredients in this formula are Schur determinants and certain integers, the *quiver coefficients*, which generalize the classical Littlewood-Richardson coefficients. Our aim in this paper is to prove a positive combinatorial formula for the quiver coefficients when the rank conditions defining the degeneracy locus are given by a permutation. In particular, this gives new expansions for Fulton's universal Schubert polynomials [F3] and the Schubert polynomials of Lascoux and Schützenberger [LS1].

Let $\mathfrak{X}$ be a smooth complex algebraic variety and let

$$(1) \quad G_1 \rightarrow \cdots \rightarrow G_{n-1} \rightarrow G_n \rightarrow F_n \rightarrow F_{n-1} \rightarrow \cdots \rightarrow F_1$$

be a sequence of vector bundles and morphisms over $\mathfrak{X}$, such that G_i and F_i have rank i for each i . For every permutation w in the symmetric group S_{n+1} there is a degeneracy locus

$$\Omega_w(G_\bullet \rightarrow F_\bullet) = \{x \in \mathfrak{X} \mid \text{rank}(G_q(x) \rightarrow F_p(x)) \leq r_w(p, q) \text{ for all } 1 \leq p, q \leq n\},$$

where $r_w(p, q)$ is the number of $i \leq p$ such that $w(i) \leq q$. The universal double Schubert polynomial $\mathfrak{S}_w(c; d)$ of Fulton gives a formula for this locus; this is a polynomial in the variables $c_i(j)$ and $d_i(j)$ for $1 \leq i \leq j \leq n$. When the codimension of $\Omega_w(G_\bullet \rightarrow F_\bullet)$ is equal to the length of w , its class $[\Omega_w]$ in the cohomology (or Chow ring) of $\mathfrak{X}$ is obtained by evaluating $\mathfrak{S}_w(c; d)$ at the Chern classes $c_i(p)$ and $d_i(q)$ of the bundles F_p and G_q , respectively. The quiver formula given in [BF] specializes to a formula for universal double Schubert polynomials:

$$\mathfrak{S}_w(c; d) = \sum_{\lambda} c_{w, \lambda}^{(n)} s_{\lambda^1}(d(2) - d(1)) \cdots s_{\lambda^n}(c(n) - d(n)) \cdots s_{\lambda^{2n-1}}(c(1) - c(2)).$$

Here the sum is over sequences of $2n - 1$ partitions $\lambda = (\lambda^1, \dots, \lambda^{2n-1})$ and each s_{λ^i} is a Schur determinant in the difference of the two alphabets in its argument. The quiver coefficients $c_{w, \lambda}^{(n)}$ can be computed by an inductive algorithm, and are conjectured to be nonnegative [BF].

Let $\text{col}(T)$ denote the column word of a semistandard Young tableau T , the word obtained by reading the entries of the columns of the tableau from bottom to top and left to right. The following theorem is our main result.

Date: November 19, 2002.

2000 *Mathematics Subject Classification.* 05E15; 14M15.

The authors were supported in part by NSF Grant DMS-0070479 (Buch), an NSF Postdoctoral Research Fellowship (Kresch), and NSF Grant DMS-0296023 (Tamvakis).

Theorem 1. *Suppose that $w \in \mathcal{S}_{n+1}$ and $\lambda = (\lambda^1, \dots, \lambda^{2n-1})$ is a sequence of partitions. Then $c_{w,\lambda}^{(n)}$ equals the number of sequences of semistandard tableaux $(T_1, \dots, T_{2n-1})$ such that the shape of T_i is conjugate to λ^i , the entries of T_i are at most $\min(i, 2n-i)$, and $\text{col}(T_1) \cdots \text{col}(T_{2n-1})$ is a reduced word for w .*

As a consequence of Theorem 1, we obtain formulas for the Schubert polynomials of Lascoux and Schützenberger, expressing them as linear combinations of products of Schur polynomials in disjoint sets of variables. The coefficients in these expansions are all quiver coefficients $c_{w,\lambda}^{(n)}$; this is true in particular for the expansion of a Schubert polynomial as a linear combination of monomials (see, e.g., [BJS, FS]).

The *Stanley symmetric functions* or stable Schubert polynomials [S] play a central role in this paper. Our main result generalizes the formula of Fomin and Greene [FG] for the expansion of these symmetric functions in the Schur basis, as well as the connection between quiver coefficients and Stanley symmetric functions obtained in [B]. In fact, we show that the universal double Schubert polynomials can be expressed as a multiplicity-free sum of products of Stanley symmetric functions (Theorem 3).

It should be noted that the formula for Schubert polynomials suggested in this paper is different from the one given in [BF]. For example, the formula from [BF, §2.3] does not make it clear that the monomial coefficients of Schubert polynomials are quiver coefficients, or even that these monomial coefficients are nonnegative.

We review the universal Schubert polynomials and Stanley symmetric functions in Section 2; in addition, we prove some required properties. In Section 3 we introduce quiver varieties and we prove Theorem 1. In the following section we apply our results to the case of ordinary Schubert polynomials. Finally, in Section 5 we use our expressions for single Schubert polynomials to obtain explicit Giambelli formulas for the classical and quantum cohomology rings of partial flag varieties.

The authors thank Sergey Fomin, Peter Magyar, and Alexander Postnikov for useful discussions.

2. PRELIMINARIES

2.1. Universal Schubert polynomials. We begin by recalling the definition of the double Schubert polynomials of Lascoux and Schützenberger [LS1, L1]. Let $X = (x_1, x_2, \dots)$ and $Y = (y_1, y_2, \dots)$ be two sequences of commuting independent variables. Given a permutation $w \in S_n$, the double Schubert polynomial $\mathfrak{S}_w(X; Y)$ is defined recursively as follows. If $w = w_0$ is the longest permutation in S_n then we set

$$\mathfrak{S}_{w_0}(X; Y) = \prod_{i+j \leq n} (x_i - y_j).$$

Otherwise we can find a simple transposition $s_i = (i, i+1) \in S_n$ such that $\ell(ws_i) = \ell(w) + 1$. Here $\ell(w)$ denotes the length of w , which is the smallest number ℓ for which w can be written as a product of ℓ simple transpositions. We then define

$$\mathfrak{S}_w(X; Y) = \partial_i(\mathfrak{S}_{ws_i}(X; Y))$$

where ∂_i is the divided difference operator given by

$$\partial_i(f) = \frac{f(x_1, \dots, x_i, x_{i+1}, \dots, x_n) - f(x_1, \dots, x_{i+1}, x_i, \dots, x_n)}{x_i - x_{i+1}}.$$

The (single) Schubert polynomial is defined by $\mathfrak{S}_w(X) = \mathfrak{S}_w(X; 0)$.

Suppose now that w is a permutation in S_{n+1} . If $u_1, \dots, u_r$ are permutations, we will write $u_1 \cdots u_r = w$ if $\ell(u_1) + \cdots + \ell(u_r) = \ell(w)$ and the product of $u_1, \dots, u_r$ is equal to w . In this case we say $u_1 \cdots u_r$ is a *reduced factorization* of w .

The *double universal Schubert polynomial* $\mathfrak{S}_w(c; d)$ of [F3] is a polynomial in the variables $c_i(j)$ and $d_i(j)$ for $1 \leq i \leq j \leq n$. For convenience we set $c_0(j) = d_0(j) = 1$ for all j and $c_i(j) = d_i(j) = 0$ if $i < 0$ or $i > j$. The classical Schubert polynomial $\mathfrak{S}_w(X)$ can be written uniquely in the form

$$\mathfrak{S}_w(X) = \sum a_{i_1, \dots, i_n} e_{i_1}(x_1) e_{i_2}(x_1, x_2) \cdots e_{i_n}(x_1, \dots, x_n)$$

where the sum is over all sequences $(i_1, \dots, i_n)$ with each $i_\alpha \leq \alpha$ and $\sum i_\alpha = \ell(w)$ [LS3]. The coefficients $a_{i_1, \dots, i_n}$ are uniquely determined integers depending on w . Define the single universal Schubert polynomial for w by

$$\mathfrak{S}_w(c) = \sum a_{i_1, \dots, i_n} c_{i_1}(1) c_{i_2}(2) \cdots c_{i_n}(n)$$

and the double universal Schubert polynomial by

$$\mathfrak{S}_w(c; d) = \sum_{u \cdot v = w} (-1)^{\ell(u)} \mathfrak{S}_{u^{-1}}(d) \mathfrak{S}_v(c).$$

Since the ordinary Schubert polynomial $\mathfrak{S}_w(X)$ does not depend on which symmetric group w belongs to, the same is true for $\mathfrak{S}_w(c; d)$.

As explained in the introduction, the double universal Schubert polynomials describe the degeneracy loci $\Omega_w(G_\bullet \rightarrow F_\bullet)$ of morphisms between vector bundles on a smooth algebraic variety $\mathfrak{X}$. The precise statement is the following result.

Theorem 2 (Fulton [F3]). *If the codimension of $\Omega_w(G_\bullet \rightarrow F_\bullet)$ is equal to $\ell(w)$ (or if this locus is empty) then the class of $\Omega_w(G_\bullet \rightarrow F_\bullet)$ in the cohomology ring of $\mathfrak{X}$ is obtained from $\mathfrak{S}_w(c; d)$ by evaluating at the Chern classes of the bundles, i.e., setting $c_i(j) = c_i(F_j)$ and $d_i(j) = c_i(G_j)$.*

We will need the following consequence of Theorem 2, which generalizes a result of Kirillov [K, (5.5)], see also [F3, (26)].

Proposition 1. *Choose $m \geq 0$ and substitute $b(j)$ for $c(j)$ and for $d(j)$ in $\mathfrak{S}_w(c; d)$ for all $j \geq m+1$. We then have*

$$\begin{aligned} \mathfrak{S}_w(c(1), \dots, c(m), b(m+1), \dots; d(1), \dots, d(m), b(m+1), \dots) \\ = \begin{cases} \mathfrak{S}_w(c; d) & \text{if } w \in S_{m+1} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Proof. If $w \in S_{m+1}$ then it follows from the definition that $\mathfrak{S}_w(c; d)$ is independent of $c_i(j)$ and $d_i(j)$ for $j \geq m+1$.

Assume that $w \in S_{n+1} \setminus S_n$ where $n > m$. We claim that $\mathfrak{S}_w(c; d)$ vanishes as soon as we set $c_i(n) = d_i(n)$. To see this, consider a variety $\mathfrak{X}$ with bundles F_j for $1 \leq j \leq n$ and G_j for $1 \leq j \leq n-1$, such that $\text{rank}(F_j) = \text{rank}(G_j) = j$, and such that all monomials of degree $\ell(w)$ in the Chern classes of these bundles are linearly independent. For example we can take a product of Grassmann varieties $\mathfrak{X} = \text{Gr}(1, N) \times \cdots \times \text{Gr}(n, N) \times \cdots \times \text{Gr}(1, N)$ for N large and use the tautological bundles.

If we set $G_n = F_n$ then we have a sequence of bundles (1) for which the map $G_n \rightarrow F_n$ is the identity and all other maps are zero. Since $r_w(n, n) = n-1$ it

follows that the locus $\Omega_w(G_\bullet \rightarrow F_\bullet)$ is empty. Theorem 2 therefore implies that $\mathfrak{S}_w(c; d)$ is zero when $c_i(n) = d_i(n) = c_i(F_n)$ for $1 \leq i \leq n$. $\square$

The next identity is due to Kirillov [K]; we include a proof for completeness.

Corollary 1 (Kirillov). *If $b_i(j)$, $c_i(j)$, and $d_i(j)$ are three sets of variables, $1 \leq i \leq j \leq n$, then we have*

$$\mathfrak{S}_w(c; d) = \sum_{u \cdot v = w} \mathfrak{S}_u(b; d) \mathfrak{S}_v(c; b).$$

Proof. Let $\mathfrak{S}(c; d)$ denote the function from permutations to polynomials which maps w to $\mathfrak{S}_w(c; d)$. Following [M1, §6] we define the product of two such functions f and g by the formula

$$(fg)(w) = \sum_{u \cdot v = w} f(u)g(v).$$

Fulton's definition of universal Schubert polynomials then says that $\mathfrak{S}(c; d) = \mathfrak{S}(0; d)\mathfrak{S}(c; 0)$, and Proposition 1 shows that $\mathfrak{S}(0; c)\mathfrak{S}(c; 0) = \mathfrak{S}(c; c) = \underline{1}$, where $\underline{1}(w) = \delta_{1, w}$. Now (6.6) of [M1] implies that also $\mathfrak{S}(c; 0)\mathfrak{S}(0; c) = \underline{1}$. We therefore obtain

$$\mathfrak{S}(c; d) = \mathfrak{S}(0; d)\mathfrak{S}(c; 0) = \mathfrak{S}(0; d)\mathfrak{S}(b; 0)\mathfrak{S}(0; b)\mathfrak{S}(c; 0) = \mathfrak{S}(b; d)\mathfrak{S}(c; b),$$

as required. $\square$

Later, we will need the following observation: Proposition 1 and Corollary 1 together imply that

$$(2) \quad \mathfrak{S}_w(c; d) = \sum_{u \cdot v = w} \mathfrak{S}_u(0, \dots, 0, c(r+1), c(r+2), \dots; d) \cdot \mathfrak{S}_v(c(1), \dots, c(r); 0)$$

and similarly,

$$(3) \quad \mathfrak{S}_w(c; d) = \sum_{u \cdot v = w} \mathfrak{S}_u(0; d(1), \dots, d(r)) \cdot \mathfrak{S}_v(c; 0, \dots, 0, d(r+1), d(r+2), \dots).$$

2.2. Symmetric functions. For each integer partition $\alpha = (\alpha_1 \geq \dots \geq \alpha_p \geq 0)$, let $|\alpha| = \sum \alpha_i$ and let α' denote the conjugate of α . Let $c = \{c_1, c_2, \dots\}$ and $d = \{d_1, d_2, \dots\}$ be ordered sets of independent variables. Define the Schur determinant

$$s_\alpha(c - d) = \det(h_{\alpha_i + j - i})_{p \times p}$$

where the elements h_k are determined by the identity of formal power series

$$\sum_{k \in \mathbb{Z}} h_k t^k = \frac{1 - d_1 t + d_2 t^2 - \dots}{1 - c_1 t + c_2 t^2 - \dots}.$$

In particular, $h_0 = 1$ and $h_k = 0$ for $k < 0$. The *supersymmetric Schur functions* $s_\alpha(X/Y)$ are obtained by setting $c_i = e_i(X)$ and $d_i = e_i(Y)$ for all i , and the usual Schur polynomials are given by the specializations

$$s_\alpha(X/Y)|_{Y=0} = s_\alpha(X) \quad \text{and} \quad s_\alpha(X/Y)|_{X=0} = (-1)^{|\alpha|} s_{\alpha'}(Y).$$

If E and F are two vector bundles with total Chern classes $c(E)$ and $c(F)$, respectively, we will denote $s_\alpha(c(E) - c(F))$ by $s_\alpha(E - F)$.

Let Λ denote the ring of symmetric functions (as in [M2]). For each permutation $w \in S_n$ there is a stable Schubert polynomial or Stanley symmetric function $F_w \in \Lambda$ which is uniquely determined by the property that

$$F_w(x_1, \dots, x_k) = \mathfrak{S}_{1^m \times w}(x_1, \dots, x_k)$$

for all $m \geq k$.¹ Here $1^m \times w \in S_{n+m}$ is the permutation which is the identity on $\{1, \dots, m\}$ and which maps j to $w(j-m) + m$ for $j > m$ (see [M1, (7.18)]). When F_w is written in the basis of Schur functions, one has

$$F_w = \sum_{\alpha: |\alpha|=\ell(w)} d_{w\alpha} s_\alpha$$

for some nonnegative integers $d_{w\alpha}$ [EG, LS2]. Fomin and Greene show that the coefficient $d_{w\alpha}$ equals the number of semistandard tableaux T of shape α' such that the column word of T is a reduced word for w [FG, Thm. 1.2].

3. QUIVER VARIETIES

3.1. Definitions. Let $E_1 \rightarrow E_2 \rightarrow \dots \rightarrow E_n$ be a sequence of vector bundles and bundle maps over a non-singular variety $\mathfrak{X}$. Given rank conditions $r = \{r_{ij}\}$ for $1 \leq i < j \leq n$ there is a *quiver variety* given by

$$\Omega_r(E_\bullet) = \{x \in \mathfrak{X} \mid \text{rank}(E_i(x) \rightarrow E_j(x)) \leq r_{ij} \ \forall i < j\}.$$

For convenience, we set $r_{ii} = \text{rank } E_i$ for all i , and we demand that the rank conditions satisfy $r_{ij} \geq \max(r_{i-1,j}, r_{i,j+1})$ and $r_{ij} + r_{i-1,j+1} \geq r_{i-1,j} + r_{i,j+1}$ for all $i \leq j$. In this case, the expected codimension of $\Omega_r(E_\bullet)$ is the number $d(r) = \sum_{i < j} (r_{i,j-1} - r_{ij})(r_{i+1,j} - r_{ij})$. The main result of [BF] states that when the quiver variety $\Omega_r(E_\bullet)$ has this codimension, its cohomology class is given by

$$(4) \quad [\Omega_r(E_\bullet)] = \sum_{\lambda} c_{\lambda}(r) s_{\lambda^1}(E_2 - E_1) \cdots s_{\lambda^{n-1}}(E_n - E_{n-1}).$$

Here the sum is over all sequences of partitions $\lambda = (\lambda^1, \dots, \lambda^{n-1})$ such that $\sum |\lambda^i| = d(r)$, and the coefficients $c_{\lambda}(r)$ are integers computed by a combinatorial algorithm which we will not reproduce here.² These coefficients are uniquely determined by the condition that (4) is true for all varieties $\mathfrak{X}$ and sequences $E_\bullet$, as well as the condition that $c_{\lambda}(r) = c_{\lambda}(r')$, where $r' = \{r'_{ij}\}$ is the set of rank conditions given by $r'_{ij} = r_{ij} + 1$ for all $i \leq j$.

The loci associated with universal Schubert polynomials are special cases of these quiver varieties. Given $w \in S_{n+1}$ we define rank conditions $r^{(n)} = \{r_{ij}^{(n)}\}$ for $1 \leq i \leq j \leq 2n$ by

$$r_{ij}^{(n)} = \begin{cases} r_w(2n+1-j, i) & \text{if } i \leq n < j \\ i & \text{if } j \leq n \\ 2n+1-j & \text{if } i \geq n+1. \end{cases}$$

Then $\Omega_w(G_\bullet \rightarrow F_\bullet)$ is identical to the quiver variety $\Omega_{r^{(n)}}(G_\bullet \rightarrow F_\bullet)$, and furthermore we have $\ell(w) = d(r)$. If we let $c_{w,\lambda}^{(n)} = c_{\lambda}(r^{(n)})$ denote the quiver coefficients

¹In Stanley's notation, the function $F_{w^{-1}}$ is assigned to w .

²Knutson, Miller, and Shimozono have recently announced that they can prove that the integers $c_{\lambda}(r)$ are nonnegative.

corresponding to this locus, it follows that

$$(5) \quad \mathfrak{S}_w(c; d) = \sum_{\lambda} c_{w, \lambda}^{(n)} s_{\lambda^1}(d(2) - d(1)) \cdots s_{\lambda^n}(c(n) - d(n)) \cdots s_{\lambda^{2n-1}}(c(1) - c(2)).$$

3.2. Proof of Theorem 1. It will be convenient to work with the element $P_w^{(n)} \in \Lambda^{\otimes 2n-1}$ defined by

$$P_w^{(n)} = \sum_{\lambda} c_{w, \lambda}^{(n)} s_{\lambda^1} \otimes \cdots \otimes s_{\lambda^{2n-1}}.$$

Theorem 1 is a consequence of Fomin and Greene's formula for stable Schubert polynomials combined with the following result.

Theorem 3. *For $w \in S_{n+1}$ we have*

$$P_w^{(n)} = \sum_{u_1 \dots u_{2n-1} = w} F_{u_1} \otimes \cdots \otimes F_{u_{2n-1}}$$

where the sum is over all reduced factorizations $w = u_1 \cdots u_{2n-1}$ such that $u_i \in S_{\min(i, 2n-i)+1}$ for each i .

Proof. Since $r_w(p, q) + m = r_{1^m \times w}(p + m, q + m)$ for $m \geq 0$, it follows that the coefficients $c_{w, \lambda}^{(n)}$ are uniquely defined by the condition that

$$(6) \quad \mathfrak{S}_{1^m \times w}(c; d) = \sum_{\lambda} c_{w, \lambda}^{(n)} s_{\lambda^1}(d(2 + m) - d(1 + m)) \cdots s_{\lambda^n}(c(n + m) - d(n + m)) \cdots s_{\lambda^{2n-1}}(c(1 + m) - c(2 + m))$$

for all $m \geq 0$ (see also [B, §4]).

Given any two integers $p \leq q$ we let $P_w^{(n)}[p, q]$ denote the sum of the terms of $P_w^{(n)}$ for which λ^i is empty when $i < p$ or $i > q$:

$$P_w^{(n)}[p, q] = \sum_{\lambda: \lambda^i = \emptyset \text{ for } i \notin [p, q]} c_{w, \lambda}^{(n)} s_{\lambda^1} \otimes \cdots \otimes s_{\lambda^{2n-1}}.$$

Lemma 1. *For any $1 < i \leq 2n - 1$ we have*

$$(7) \quad P_w^{(n)} = \sum_{u \cdot v = w} P_u^{(n)}[1, i - 1] \cdot P_v^{(n)}[i, 2n - 1].$$

Proof. We will do the case $i \leq n$; the other one is similar. For $f = \sum c_{\lambda} s_{\lambda^1} \otimes \cdots \otimes s_{\lambda^{2n-1}} \in \Lambda^{\otimes 2n-1}$, we set

$$f(c; d) = \sum_{\lambda} c_{\lambda} s_{\lambda^1}(d(2) - d(1)) \cdots s_{\lambda^n}(c(N) - d(N)) \cdots s_{\lambda^{2n-1}}(c(1) - c(2)).$$

Equation (6) implies that $P_w^{(n)} \in \Lambda^{\otimes 2n-1}$ is the unique element satisfying that $(1^{\otimes m} \otimes P_w^{(n)} \otimes 1^{\otimes m})(c; d) = \mathfrak{S}_{1^m \times w}(c; d)$ for all m . This uniqueness is preserved even if we set $d(i + m) = 0$. The right hand side of the identity (7) satisfies this by equation (3) applied to $1^m \times w$. $\square$

Lemma 2. *For $1 \leq i \leq 2n - 1$ we have*

$$P_w^{(n)}[i, i] = \begin{cases} 1^{\otimes i-1} \otimes F_w \otimes 1^{\otimes 2n-1-i} & \text{if } w \in S_{m+1}, m = \min(i, 2n - i) \\ 0 & \text{otherwise.} \end{cases}$$

Proof. If $w \notin S_{m+1}$ then this follows from Proposition 1, so assume $w \in S_{m+1}$. For simplicity we will furthermore assume that $m = i$. It is proved in [B, §4] that $P_w^{(m)}[m, m] = 1^{\otimes m-1} \otimes F_w \otimes 1^{\otimes m-1}$. Let $\Phi : \Lambda^{\otimes 2m-1} \rightarrow \Lambda^{\otimes 2n-1}$ be the linear map given by

$$\Phi(s_{\lambda^1} \otimes \cdots \otimes s_{\lambda^{2m-1}}) = s_{\lambda^1} \otimes \cdots \otimes s_{\lambda^{m-1}} \otimes \Delta^{2n-2m}(s_{\lambda^m}) \otimes s_{\lambda^{m+1}} \otimes \cdots \otimes s_{\lambda^{2m-1}}$$

where $\Delta^{2n-2m} : \Lambda \rightarrow \Lambda^{\otimes 2n-2m+1}$ denotes the $2n - 2m$ fold coproduct, that is

$$\Delta^{2n-2m}(s_{\lambda^m}) = \sum_{\tau_1, \dots, \tau_{2n-2m+1}} c_{\tau_1, \dots, \tau_{2n-2m+1}}^{\lambda^m} s_{\tau_1} \otimes \cdots \otimes s_{\tau_{2n-2m+1}}$$

where $c_{\tau_1, \dots, \tau_{2n-2m+1}}^{\lambda^m}$ is the coefficient of s_{λ^m} in the product $s_{\tau_1} s_{\tau_2} \cdots s_{\tau_{2n-2m+1}}$. For the definition of the locus $\Omega_w(G_\bullet \rightarrow F_\bullet)$, the bundles F_i and G_i for $i \geq m+1$ are inessential in the sense of [BF, §4]. Equation (4.2) of loc. cit. therefore implies that $\Phi(P_w^{(m)}) = P_w^{(n)}$. Now the result follows from the identity $P_w^{(n)}[m, 2n-m] = \Phi(P_w^{(m)}[m, m]) = 1^{\otimes m-1} \otimes \Delta^{2n-2m}(F_w) \otimes 1^{\otimes m-1}$. $\square$

Theorem 3 follows immediately from lemmas 1 and 2. $\square$

Example 1. For the permutation $w = s_2 s_1 = 312$ in S_3 , the sequences of tableaux which satisfy the conditions of Theorem 1 are

$$(\emptyset, \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \emptyset) \quad \text{and} \quad (\emptyset, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \end{bmatrix}).$$

It follows that

$$\begin{aligned} \mathfrak{S}_{312}(c; d) &= s_2(c(2) - d(2)) + s_1(c(2) - d(2))s_1(c(1) - c(2)) \\ &= c_1(1)c_1(2) - c_1(1)d_1(2) - c_2(2) + d_2(2). \end{aligned}$$

In [BF], a conjectural combinatorial rule for general quiver coefficients $c_\lambda(r)$ was given. Although this rule was also stated in terms of sequences of semistandard tableaux satisfying certain conditions, it is different from Theorem 1 in the case of universal Schubert polynomials. It would be interesting to find a bijection that establishes the equivalence of these two rules.

3.3. Skipping bundles. A permutation w has a descent position at i if $w(i) > w(i+1)$. We say that a sequence $\{a_k\} : a_1 < \cdots < a_p$ of integers is *compatible with* w if all descent positions of w are contained in $\{a_k\}$. Suppose that $w \in S_{n+1}$ and let $1 \leq a_1 < a_2 < \cdots < a_p \leq n$ and $1 \leq b_1 < b_2 < \cdots < b_q \leq n$ be two sequences compatible with w and w^{-1} , respectively.

We let $E_\bullet$ denote the subsequence

$$G_{b_1} \rightarrow G_{b_2} \rightarrow \cdots \rightarrow G_{b_q} \rightarrow F_{a_p} \rightarrow \cdots \rightarrow F_{a_2} \rightarrow F_{a_1},$$

and define rank conditions $\tilde{r}^{(n)} = \{\tilde{r}_{ij}^{(n)}\}$ for $1 \leq i \leq j \leq p+q$ by

$$\tilde{r}_{ij}^{(n)} = \begin{cases} r_w(a_{p+q+1-j}, b_i) & \text{if } i \leq q < j \\ b_i & \text{if } j \leq q \\ a_{p+q+1-j} & \text{if } i \geq q+1 \end{cases}$$

Then the expected codimension of the locus $\Omega_{\tilde{r}^{(n)}}(E_\bullet)$ is equal to $\ell(w)$. However, in general this locus may contain $\Omega_w(G_\bullet \rightarrow F_\bullet)$ as a proper closed subset. We will need the following criterion for equality (see also the remarks in [F3, §3] and Exercise 10 of [F2, §10].)

Lemma 3. *Suppose that the map $G_{i-1} \rightarrow G_i$ is injective for $i \notin \{b_k\}$ and the map $F_i \rightarrow F_{i-1}$ is surjective for $i \notin \{a_k\}$. Then $\Omega_{\tilde{r}(n)}(E_\bullet) = \Omega_w(G_\bullet \rightarrow F_\bullet)$ as subschemes of $\mathfrak{X}$.*

Proof. Let $1 \leq i, j \leq n$ be given. If i is not a descent position for w then either $w(i) \leq j$ or $w(i+1) > j$. In the first case this implies that $r_w(i, j) = r_w(i-1, j) + 1$ so the condition $\text{rank}(G_j \rightarrow F_i) \leq r_w(i, j)$ follows from $\text{rank}(G_j \rightarrow F_{i-1}) \leq r_w(i-1, j)$ because the map $F_i \rightarrow F_{i-1}$ is surjective. In the second case we have $r_w(i, j) = r_w(i+1, j)$ so the rank condition on $G_j \rightarrow F_i$ follows from the one on $G_j \rightarrow F_{i+1}$. A similar argument works if w^{-1} does not have a descent at position j . We conclude that the locus $\Omega_w(G_\bullet \rightarrow F_\bullet)$ does not change if the bundles G_j for $j \notin \{b_k\}$ and F_i for $i \notin \{a_k\}$ are disregarded. $\square$

Corollary 2. *Let $w \in S_{n+1}$ and $\{a_k\}$ and $\{b_k\}$ be as above. Then we have*

$$[\Omega_{\tilde{r}(n)}(E_\bullet)] = \sum_{\mu} \tilde{c}_{w,\mu}^{(n)} s_{\mu^1}(G_{b_2} - G_{b_1}) \cdots s_{\mu^q}(F_{a_p} - G_{b_q}) \cdots s_{\mu^{p+q-1}}(F_{a_1} - F_{a_2})$$

with coefficients $\tilde{c}_{w,\mu}^{(n)} = c_{w,\lambda}^{(n)}$ where the sequence $\lambda = (\lambda^1, \dots, \lambda^{2n-1})$ is given by

$$\lambda^i = \begin{cases} \mu^k & \text{if } i = b_k \\ \mu^{p+q-k} & \text{if } i = 2n - a_k \\ \emptyset & \text{otherwise.} \end{cases}$$

Proof. We may assume that $F_i = F_{i-1} \oplus \mathbb{C}$ when $i \notin \{a_k\}$ and $G_j = G_{j-1} \oplus \mathbb{C}$ when $j \notin \{b_k\}$. In this case, notice that $s_\alpha(F_i - F_{i-1})$ is non-zero only when α is the empty partition or when $i = a_k$ for some k . Similarly $s_\alpha(G_{i-1} - G_i)$ is zero unless α is empty or $i = b_k$ for some k . The result therefore follows from Lemma 3 and equation (5). $\square$

4. SCHUBERT POLYNOMIALS

4.1. Degeneracy loci. In this section, we will interpret the previous results for ordinary double Schubert polynomials. Let V be a vector bundle of rank n and let

$$G_1 \subset G_2 \subset \cdots \subset G_{n-1} \subset V \twoheadrightarrow F_{n-1} \twoheadrightarrow \cdots \twoheadrightarrow F_2 \twoheadrightarrow F_1$$

be a complete flag followed by a dual complete flag of V . If $w \in S_n$ then Fulton has proved [F1] that

$$[\Omega_w(G_\bullet \rightarrow F_\bullet)] = \mathfrak{S}_w(x_1, \dots, x_n; y_1, \dots, y_n)$$

where $x_i = c_1(\ker(F_i \rightarrow F_{i-1}))$, $y_i = c_1(G_i/G_{i-1})$, and $\mathfrak{S}_w(X; Y)$ is the double Schubert polynomial of Lascoux and Schützenberger.

Set $G'_i = V/G_i$ and $F'_i = \ker(V \rightarrow F_i)$. Then we have a sequence

$$F'_{n-1} \subset \cdots \subset F'_1 \subset V \twoheadrightarrow G'_1 \twoheadrightarrow \cdots \twoheadrightarrow G'_{n-1}$$

and it is easy to check that $\Omega_w(G_\bullet \rightarrow F_\bullet) = \Omega_{w_0 w^{-1} w_0}(F'_\bullet \rightarrow G'_\bullet)$ as subschemes of $\mathfrak{X}$, where w_0 is the longest permutation in S_n .

Let $1 \leq a_1 < \cdots < a_p \leq n-1$ and $0 \leq b_1 < \cdots < b_q \leq n-1$ be two sequences compatible with w and w^{-1} , respectively. Then by applying section 3.3 to the

subsequence $F'_{a_p} \rightarrow \cdots \rightarrow F'_{a_1} \rightarrow G'_{b_1} \rightarrow \cdots \rightarrow G'_{b_q}$ we obtain

$$[\Omega_w(G_\bullet \rightarrow F_\bullet)] = \sum_{\mu} \tilde{c}_{w_0 w^{-1} w_0, \mu}^{(n-1)} s_{\mu^1}(F'_{a_{p-1}} - F'_{a_p}) \cdots s_{\mu^p}(G'_{b_1} - F'_{a_1}) \cdots s_{\mu^{p+q-1}}(G'_{b_q} - G'_{b_{q-1}}).$$

Set $a_0 = b_0 = 0$. If we let $X_i = \{x_{a_{i-1}+1}, \dots, x_{a_i}\}$ denote the Chern roots of $\ker(F_{a_i} \rightarrow F_{a_{i-1}})$ and $Y_i = \{y_{b_{i-1}+1}, \dots, y_{b_i}\}$ be the Chern roots of $G_{b_i}/G_{b_{i-1}}$ then the previous equality can be written as

$$(8) \quad \mathfrak{S}_w(X; Y) = \sum_{\mu} \tilde{c}_{w_0 w^{-1} w_0, \mu}^{(n-1)} s_{\mu^1}(X_p) \cdots s_{\mu^p}(X_1/Y_1) \cdots s_{\mu^{p+q-1}}(0/Y_q).$$

This equation is true in the cohomology ring $H^*(\mathfrak{X}; \mathbb{Z})$, in which there are relations between the variables x_i and y_i (including e.g. the relations $e_j(x_1, \dots, x_n) = c_j(V) = e_j(y_1, \dots, y_n)$ for $1 \leq j \leq n$). We claim, however, that (8) holds as an identity of polynomials in independent variables. For this, notice that the identity is independent of n , i.e. the coefficient $\tilde{c}_{w_0 w^{-1} w_0, \mu}^{(n-1)}$ does not change when n is replaced with $n+1$ and w_0 with the longest element in S_{n+1} . If we choose n sufficiently large, we can construct a variety $\mathfrak{X}$ on which (8) is true, and where all monomials in the variables x_i and y_i of total degree at most $\ell(w)$ are linearly independent, which establishes the claim.

4.2. Splitting Schubert polynomials. We continue by reformulating equation (8) to obtain a more natural expression for double Schubert polynomials.

It follows from (8) together with the main result of [B] that $F_{w_0 w^{-1} w_0} = F_w$. Therefore, the coefficient $d_{w\alpha}$ of the Schur expansion of F_w is equal to the number of semistandard tableaux of shape α' such that the column word is a reduced word for $w_0 w^{-1} w_0$. Notice that if $e = (e_1, e_2, \dots, e_\ell)$ is a reduced word for $w_0 w^{-1} w_0$ then $\tilde{e} = (n+1-e_\ell, \dots, n+1-e_1)$ is a reduced word for w . Furthermore, if e is the column word of a tableau of shape α' , then $\tilde{e}$ is the column word of a skew tableau whose shape is the 180 degree rotation of α' . If we denote this rotated shape by $\tilde{\alpha}$ then we conclude that $d_{w\alpha}$ is also equal to the number of skew tableaux of shape $\tilde{\alpha}$ such that the column word is a reduced word for w . Theorem 3 now implies the following variation of our main theorem:

Theorem 1'. *If $w \in S_{n+1}$ then the coefficient $c_{w, \lambda}^{(n)}$ equals the number of sequences of semistandard skew tableaux $(T_1, \dots, T_{2n-1})$ such that T_i has shape $\tilde{\lambda}^i$, the entries of T_i are at most $\min(i, 2n-i)$, and $\text{col}(T_1) \cdots \text{col}(T_{2n-1})$ is a reduced word for w .*

We say that a sequence of tableaux $(T_1, \dots, T_r)$ is *strictly bounded below* by an integer sequence $(a_1, \dots, a_r)$ if the entries of T_i are strictly greater than a_i , for each i .

Theorem 4. *Let $w \in S_n$ and let $1 \leq a_1 < \cdots < a_p$ and $0 \leq b_1 < \cdots < b_q$ be two sequences compatible with w and w^{-1} , respectively. Then we have*

$$(9) \quad \mathfrak{S}_w(X; Y) = \sum_{\lambda} c_{\lambda} s_{\lambda^1}(0/Y_q) \cdots s_{\lambda^q}(X_1/Y_1) \cdots s_{\lambda^{p+q-1}}(X_p)$$

where $X_i = \{x_{a_{i-1}+1}, \dots, x_{a_i}\}$ and $Y_i = \{y_{b_{i-1}+1}, \dots, y_{b_i}\}$ and the sum is over all sequences of partitions $\lambda = (\lambda^1, \dots, \lambda^{p+q-1})$. Each c_{λ} is a quiver coefficient, equal to the number of sequences of semistandard tableaux $(T_1, \dots, T_{p+q-1})$ strictly

bounded below by $(b_{q-1}, \dots, b_1, 0, a_1, a_2, \dots, a_{p-1})$, such that the shape of T_i is conjugate to λ^i and $\text{col}(T_1) \cdots \text{col}(T_{p+q-1})$ is a reduced word for w .

Proof. This follows from equation (8) together with Theorem 1' applied to $w_0 w^{-1} w_0$. To translate between the sequences of skew tableaux in Theorem 1' and the sequences in the present theorem, simply rotate a whole sequence of skew tableaux by 180 degrees (this means invert the order of the sequence, and turn each skew tableau on its head). Then replace each entry e with $n + 1 - e$.

A purely algebraic proof of the splitting formula (9) is also possible. This requires versions of the results of Section 2.1 for ordinary Schubert polynomials, available e.g. from [L2] and [FK], together with [FG, Thm. 1.2]. However, some geometric reasoning is needed to interpret the c_λ as quiver coefficients. $\square$

Notice that if one takes $b_1 = 0$ in Theorem 4 then the set of variables Y_1 is empty, so equation (9) contains only signed products of single Schur polynomials. Observe also that a factor $s_{\lambda^i}(0/Y_k)$ in (9) will vanish if λ^i has more than $b_k - b_{k-1}$ columns and that $s_{\lambda^i}(X_k)$ vanishes if λ^i has more than $a_k - a_{k-1}$ rows. Therefore equation (9) uses only a subset of the quiver coefficients of Theorem 4.

Example 2. Let $w = 321$ be the longest element in S_3 , and choose the sequences $\{a_i\} = \{b_i\} = \{1 < 2\}$. Then the four sequences of tableaux satisfying the conditions of Theorem 4 are

$$(\boxed{2}, \boxed{1}, \boxed{2}), \quad (\boxed{2}, \boxed{12}, \emptyset), \quad (\emptyset, \boxed{\frac{1}{2}}, \boxed{2}), \quad \text{and} \quad (\emptyset, \boxed{\frac{1}{2}}, \emptyset),$$

all of which give nonvanishing terms and correspond to the reduced word $s_2 s_1 s_2$. We thus have

$$\begin{aligned} \mathfrak{S}_{321}(X; Y) &= \\ s_1(0/y_2)s_1(x_1/y_1)s_1(x_2) &+ s_1(0/y_2)s_{1,1}(x_1/y_1) + s_2(x_1/y_1)s_1(x_2) + s_{2,1}(x_1/y_1) \\ &= -y_2(x_1 - y_1)x_2 - y_2(y_1^2 - x_1y_1) + (x_1^2 - x_1y_1)x_2 + (x_1y_1^2 - x_1^2y_1) \\ &= (x_1 - y_1)(x_1 - y_2)(x_2 - y_1). \end{aligned}$$

Corollary 3. Suppose that $w \in S_n$ is a permutation compatible with the sequence $a_1 < \cdots < a_p$. Then we have

$$\mathfrak{S}_w(X) = \sum_{\lambda} c_{\lambda} s_{\lambda^1}(X_1) \cdots s_{\lambda^p}(X_p)$$

where $X_i = \{x_{a_{i-1}+1}, \dots, x_{a_i}\}$ and the sum is over all sequences of partitions $\lambda = (\lambda^1, \dots, \lambda^p)$. Each c_{λ} is a quiver coefficient, equal to the number of sequences of semistandard tableaux $(T_1, \dots, T_p)$ strictly bounded below by $(0, a_1, a_2, \dots, a_{p-1})$, such that the shape of T_i is conjugate to λ^i and $\text{col}(T_1) \cdots \text{col}(T_p)$ is a reduced word for w .

Example 3. Consider the permutation $w = s_1 s_2 s_1 s_3 s_4 s_3 = 32541$ in S_5 , with descent positions at 1, 3, and 4. The two sequences of tableaux satisfying the conditions of Corollary 3 which give nonvanishing terms are

$$(\boxed{\frac{1}{2}}, \boxed{23}, \boxed{4}) \quad \text{and} \quad (\boxed{\frac{1}{2}}, \boxed{\frac{2}{4}}, \boxed{4}).$$

The reduced words for w corresponding to these sequences are $s_4 s_2 s_1 s_2 s_3 s_4$ and $s_2 s_1 s_4 s_2 s_3 s_4$, respectively. It follows that

$$\begin{aligned}\mathfrak{S}_{32541}(X) &= s_3(x_1)s_{1,1}(x_2, x_3)s_1(x_4) + s_2(x_1)s_{2,1}(x_2, x_3)s_1(x_4) \\ &= x_1^3 \cdot x_2 x_3 \cdot x_4 + x_1^2(x_2^2 x_3 + x_3^2 x_2)x_4.\end{aligned}$$

The special case of Corollary 3 with $a_k = k$ gives a formula for the coefficient of each monomial in $\mathfrak{S}_w(X)$, which is equivalent to that of [BJS, Thm. 1.1]. We deduce that these monomial coefficients are quiver coefficients. The same conclusion holds for double Schubert polynomials:

Corollary 4. *Let $w \in S_n$ and let $x^u y^v = x_1^{u_1} \cdots x_{n-1}^{u_{n-1}} y_1^{v_1} \cdots y_{n-1}^{v_{n-1}}$ be a monomial of total degree $\ell(w)$. Set $g_i = \sum_{k=n-i}^{n-1} v_k$ and $f_i = g_{n-1} + \sum_{k=1}^i u_k$. Then the coefficient of $x^u y^v$ in the double Schubert polynomial $\mathfrak{S}_w(X; Y)$ is equal to $(-1)^{g_{n-1}}$ times the number of reduced words $(e_1, \dots, e_{\ell(w)})$ for w such that $n-i \leq e_{g_{i-1}+1} < \cdots < e_{g_i}$ and $e_{f_{i-1}+1} > \cdots > e_{f_i} \geq i$ for all $1 \leq i \leq n-1$.*

Proof. Take $\{a_k\} = \{1, 2, \dots, n-1\}$ and $\{b_k\} = \{0, 1, 2, \dots, n-1\}$ in Theorem 4 and notice that a Schur polynomial $s_\alpha(0/y_i)$ is non-zero only if $\alpha = (1^{v_i})$ is a single column with v_i boxes, in which case $s_\alpha(0/y_i) = (-y_i)^{v_i}$. Similarly $s_\alpha(x_i)$ is equal to $x_i^{u_i}$ if $\alpha = (u_i)$ is a single row with u_i boxes, and is zero otherwise. The reduced words of the corollary are exactly those that form a sequence of tableaux on the conjugates of these shapes and strictly bounded below by $(n-2, n-3, \dots, 1, 0, 0, 1, \dots, n-2)$. $\square$

Remark. Corollary 3 implies that the single Schubert polynomial $\mathfrak{S}_w(X)$ is the character of a $\mathrm{GL}(a_1, \mathbb{C}) \times \mathrm{GL}(a_2 - a_1, \mathbb{C}) \times \cdots \times \mathrm{GL}(a_p - a_{p-1}, \mathbb{C})$ module. Our expression for the coefficients c_λ gives an isotypic decomposition of this module. This is closely related to a theorem of Kraskiewicz and Pragacz [KP], which shows that $\mathfrak{S}_w(X)$ is a character of a Borel module. P. Magyar reports that the former property may be deduced from the latter (private communication).

5. GIAMBELLI FORMULAS

In this final section, we observe that Corollaries 2 and 3 give explicit, non-recursive answers to the *Giambelli problem* for the classical and quantum cohomology of partial flag manifolds.

Suppose that V is a complex vector space of dimension n and choose integers $0 = a_0 < a_1 < \cdots < a_p < a_{p+1} = n$. Let $\mathfrak{X}$ be the partial flag variety which parametrizes quotients

$$(10) \quad V \twoheadrightarrow F_{a_p} \twoheadrightarrow \cdots \twoheadrightarrow F_{a_1}$$

with $\mathrm{rank}(F_{a_k}) = a_k$ for each k . We will also use (10) to denote the tautological sequence of quotient bundles over $\mathfrak{X}$, and define $Q_k = \mathrm{Ker}(F_{a_k} \rightarrow F_{a_{k-1}})$, for $1 \leq k \leq p+1$. According to Borel [Bo], the cohomology ring $H^*(\mathfrak{X}; \mathbb{Z})$ is presented as the polynomial ring in the Chern classes $c_i(Q_k)$ for all i and k , modulo the relation

$$c(Q_1)c(Q_2) \cdots c(Q_{p+1}) = 1.$$

Fix a complete flag $G_\bullet$ of subspaces of V , and let $S(a)$ denote the subset of S_n consisting of permutations w compatible with $\{a_k\}$. For each $w \in S(a)$, there is a Schubert variety $\Omega_w \subset \mathfrak{X}$, defined as the locus of $x \in \mathfrak{X}$ such that

$$\mathrm{rank}(G_j(x) \rightarrow F_i(x)) \leq r_w(i, j), \quad \text{for } i \in \{a_1, \dots, a_p\} \text{ and } 1 \leq j \leq n.$$

The *Schubert classes* $[\Omega_w] \in H^{2\ell(w)}(\mathfrak{X}; \mathbb{Z})$ for $w \in S(a)$ form a natural ‘geometric basis’ for the cohomology ring of $\mathfrak{X}$. The following Giambelli formula, which is a direct consequence of Corollary 3, writes these classes as polynomials in the ‘algebraic generators’ for $H^*(\mathfrak{X}; \mathbb{Z})$ given by the $c_i(Q_k)$.

Giambelli I. *We have*

$$(11) \quad [\Omega_w] = \sum_{\lambda} c_{\lambda} s_{\lambda^1}(Q_1) \cdots s_{\lambda^p}(Q_p),$$

where the sum is over sequences of partitions $\lambda = (\lambda^1, \dots, \lambda^p)$ and c_{λ} is the quiver coefficient of Corollary 3.

Alternatively, we can use Corollary 2 to express the Schubert class $[\Omega_w]$ as a polynomial in the special Schubert classes $c_i(F_{a_k})$. Notice that the two approaches are identical for Grassmannians.

Giambelli II. *We have*

$$(12) \quad [\Omega_w] = \sum_{\nu} \tilde{c}_{\nu} s_{\nu^p}(F_{a_p}) \cdots s_{\nu^1}(F_{a_1} - F_{a_2}),$$

where the sum is over sequences of partitions $\nu = (\nu^1, \dots, \nu^p)$ and $\tilde{c}_{\nu} = c_{w, \lambda}^{(n)}$, where $\lambda = (\lambda^1, \dots, \lambda^{2n-1})$ is given by

$$\lambda^i = \begin{cases} \nu^k & \text{if } i = 2n - a_{p+1-k} \\ \emptyset & \text{otherwise.} \end{cases}$$

When comparing the above two Giambelli formulas, recall that the quiver coefficients c_{λ} which appear in (11) correspond to the permutation $w_0 w^{-1} w_0$. The equivalence of (11) and (12) can also be checked directly, by using the Chern class identities $c(Q_k) = c(F_{a_k} - F_{a_{k-1}})$.

Following [FGP] and [C-F, §3.2], we recall that equation (12) can be used to obtain a quantum Giambelli formula which holds in the small quantum cohomology ring $QH^*(\mathfrak{X})$. For this, one simply replaces all the special Schubert classes which appear in the Schur determinants $s_{\nu^k}(F_{a_k} - F_{a_{k+1}})$ in (12) with the corresponding quantum classes, as in loc. cit. (compare also with [F3, Prop. 4.3]).

REFERENCES

- [BJS] S. Billey, W. Jockusch and R. P. Stanley : *Some combinatorial properties of Schubert polynomials*, J. Algebraic Combin. **2** (1993), no. 4, 345–374.
- [Bo] A. Borel : *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Ann. of Math. **57** (1953), 115–207.
- [B] A. S. Buch : *Stanley symmetric functions and quiver varieties*, J. Algebra **235** (2001), no. 1, 243–260.
- [BF] A. S. Buch and W. Fulton : *Chern class formulas for quiver varieties*, Invent. Math. **135** (1999), no. 3, 665–687.
- [C-F] I. Ciocan-Fontanine : *On quantum cohomology rings of partial flag varieties*, Duke Math. J. **98** (1999), no. 3, 485–524.
- [EG] M. Edelman and C. Greene : *Balanced tableaux*, Adv. Math. **63** (1987), 42–99.
- [FGP] S. Fomin, S. Gelfand and A. Postnikov : *Quantum Schubert polynomials*, J. Amer. Math. Soc. **10** (1997), 565–596.
- [FG] S. Fomin and C. Greene : *Noncommutative Schur functions and their applications*, Discrete Math. **193** (1998), no. 1–3, 179–200.
- [FK] S. Fomin and A. N. Kirillov : *The Yang-Baxter equation, symmetric functions, and Schubert polynomials*, Discrete Math. **153** (1996), no. 1–3, 123–143.

- [FS] S. Fomin and R. P. Stanley : *Schubert polynomials and the nilCoxeter algebra*, Adv. Math. **103** (1994), 196–207.
- [F1] W. Fulton : *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, Duke Math. J. **65** (1992), no. 3, 381–420.
- [F2] W. Fulton : *Young Tableaux*, L.M.S. Student Texts **35**, Cambridge University Press, Cambridge, 1997.
- [F3] W. Fulton : *Universal Schubert polynomials*, Duke Math. J. **96** (1999), no. 3, 575–594.
- [K] A. N. Kirillov : *Cauchy identities for universal Schubert polynomials*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **283** (2001), 123–139.
- [KP] W. Kraskiewicz and P. Pragacz : *Foncteurs de Schubert*, C. R. Acad. Sci. Paris Sér. I Math. **304** (1987), 209–211.
- [L1] A. Lascoux : *Classes de Chern des variétés de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), no. 5, 393–398.
- [L2] A. Lascoux : *Notes on interpolation in one and several variables*, available at <http://phalanstere.univ-mlv.fr/~al/>
- [LS1] A. Lascoux and M.-P. Schützenberger : *Polynômes de Schubert*, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982), 447–450.
- [LS2] A. Lascoux and M.-P. Schützenberger : *Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), 629–633.
- [LS3] A. Lascoux and M.-P. Schützenberger : *Funtorialité des polynomes de Schubert*, Contemporary Math **88** (1989), 585–598.
- [M1] I. G. Macdonald : *Notes on Schubert polynomials*, Publ. LACIM **6**, Univ. de Québec à Montréal, Montréal, 1991.
- [M2] I. G. Macdonald : *Symmetric Functions and Hall Polynomials*, Second edition, Clarendon Press, Oxford, 1995.
- [S] R. Stanley : *On the number of reduced decompositions of elements of Coxeter groups*, European J. Combin. **5** (1984), 359–372.

MATEMATISK INSTITUT, AARHUS UNIVERSITET, NY MUNKEGADE, 8000 ÅRHUS C, DENMARK
E-mail address: `abuch@imf.au.dk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, 209 SOUTH 33RD STREET,
 PHILADELPHIA, PA 19104-6395, USA
E-mail address: `kresch@math.upenn.edu`

DEPARTMENT OF MATHEMATICS, BRANDEIS UNIVERSITY - MS 050, P. O. BOX 9110, WALTHAM,
 MA 02454-9110, USA
E-mail address: `harryt@brandeis.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 525 EAST UNIVERSITY AVE., ANN
 ARBOR, MI 48109-1109, USA
E-mail address: `ayong@umich.edu`